

Centre de Physique Théorique* - CNRS - Luminy, Case 907 F-13288
Marseille Cedex 9 - France

EXISTENCE OF LONG-RANGE ORDER IN QUASI-TWO-DIMENSIONAL HUBBARD MODEL

A. BELKASRI¹ and J.L. RICHARD

Abstract

In recent work of Monthoux and Pines [1] and also in Rice et *al.*'s work [2], quasi-averages like $\langle c_{k\uparrow} c_{-k\downarrow} \rangle$ were considered even in the case of a dimension less or equal two. But it is well known from the old work of Hohenberg [3] that these quasi-averages are zero at $T \neq 0$ in case of 1 and 2 dimensions. In this communication we apply the result of Hohenberg to the Hubbard model and prove that in the case of quasi-two-dimension, the inequality of Bogoliubov is not in contradiction with having $\langle c_{k\uparrow} c_{-k\downarrow} \rangle \neq 0$ (at $T \neq 0$) even for a system of three layers.

Key-Words : Strongly Correlated Systems, Off Diagonal LRO, HTC superconductivity.

July 1994
CPT-94/P.3025

anonymous ftp or gopher: cpt.univ-mrs.fr

* Unité Propre de Recherche 7061

¹and Université d'Aix-Marseille II

The models which are generally used to describe the electronic and magnetic properties of oxides HT_c superconductors are based on the Hubbard model. The HT_c oxides present a layered structure and copper-oxide planes are the main structural elements of these materials. Therefore the relevant properties of these oxides refer to a two dimensional systems.

We know from the work of Hohenberg [3] that a two dimensional Fermi (or Bose) system can not exhibit a Long-Range Order (LRO) for $T \neq 0$. Nevertheless recently many authors [1, 2, 4] considered a LRO for models derived from the Hubbard model even in two-dimensions.

In BCS theory superconductivity is associated with anomalous averages $\langle c_{k\uparrow}c_{-k\downarrow} \rangle$ which are non zero by virtue of a broken symmetry (the conservation of particle number). Indeed, the key feature of BCS theory is Cooper-pair condensation. The pair of states $(k \uparrow, -k \downarrow)$ is occupied coherently. The Cooper-pair amplitude, $\langle c_{k\uparrow}c_{-k\downarrow} \rangle$, which is zero in the normal state, becomes finite below T_c .

In what follows, we repeat the proof of Hohenberg [3] for the case of Hubbard model and show that in quasi-two-dimensional case a LRO may exist.

1 LRO in Hubbard model

1.1 Hubbard model of fermions

In this section we will examine the ordering for fermions in the frame work of a quasi-two-dimensional Hubbard model. Let us consider the following Hamiltonian $H = H_t + H_U$, with

$$H_t = \sum_{ij} t_{ij} c_{i\sigma}^+ c_{j\sigma} \quad (1)$$

and

$$H_U = \sum_{i,j;\sigma,\sigma'} U_{ij} n_{i\sigma} n_{j\sigma'} \quad (2)$$

where $c_{i\sigma}$, $c_{i\sigma}^+$ are the destruction and creation operators, respectively, and $n_{i\sigma} = c_{i\sigma}^+ c_{i\sigma}$ with σ is the spin index. The hopping amplitude t_{ij} is nonzero only for (i, j) nearest-neighbors, and we have

$$t_{ij} = \begin{cases} -t & \text{for } i - j = \pm e_x \text{ or } \pm e_y \\ -t' & \text{for } i - j = \pm e_z \\ 0 & \text{otherwise} \end{cases} \quad (3)$$

with (e_x, e_y, e_z) the unit vectors for the three directions of the lattice. The amplitude of the interaction U_{ij} is effective only in directions (e_x, e_y) .

Let us define the charge density operator

$$\rho(i) = \sum_{\sigma} c_{i\sigma}^+ c_{i\sigma} \quad (4)$$

Its q^{th} Fourier component is given by

$$\rho_q = \sum_{k,\sigma} c_{k,\sigma}^+ c_{k+q,\sigma} \quad (5)$$

Since we will examine the existence or not of the quasi-averages $\langle c_{k\uparrow} c_{-k\downarrow} \rangle$ in one or more regions of k , we introduce the following order parameter

$$\Delta \equiv \frac{1}{N} \sum_k S(k) \langle c_{k\uparrow} c_{-k\downarrow} \rangle \quad (6)$$

where the “smearing function” $S(k)$ is arbitrary (a Gaussian, for instance) but has the properties

$$S(0) = 1, \quad \frac{1}{N} \sum_k S(k) < \infty \quad (7)$$

where N is the volume.

Now we will use the Bogoliubov inequality [3] which is

$$\langle \{A, A^+\} \rangle \langle [C, H], C^+ \rangle \geq 2k_B T |\langle [C, A] \rangle|^2 \quad (8)$$

where $\{.,.\}$ is an anti-commutator, $[.,.]$ is a commutator, H is any Hamiltonian, C, A are operators and finally $\langle \dots \rangle$ indicates a statistical average on a grand canonical ensemble

$$\langle X \rangle = \text{Tr} (X e^{-\beta H}) / \text{Tr} e^{-\beta H} \quad (9)$$

straightforward algebraic calculation leads to

$$[\rho_q, H_t], \rho_q^+ = -2 \sum_{i,j,\sigma} t_{ij} [1 - \cos q(i-j)] c_{i\sigma}^+ c_{i\sigma} \quad (10)$$

In Fourier transform we obtain

$$\begin{aligned} [\rho_q, H_t], \rho_q^+ = & 4t \sum_{k\sigma} \{ (1 - \cos q_x) \cos k_x + \\ & + (1 - \cos q_y) \cos k_y + \eta (1 - \cos q_z) \cos k_z \} n_{k\sigma} \end{aligned} \quad (11)$$

with $\eta = t'/t$. Since H_U commute with ρ_q , it is easy to get the following inequalities

$$0 < \langle [\rho_q, H], \rho_q^+ \rangle \leq 4t \{ (1 - \cos q_x) + (1 - \cos q_y) + \eta (1 - \cos q_z) \} \sum_k \langle n_k \rangle \quad (12)$$

and we have

$$\sum_k \langle n_k \rangle = nN \quad (13)$$

where n is the density of particles.

If in Bogoliubov inequality (Eq. (8)), we set

$$C = \rho_q \quad \text{and} \quad A = \frac{1}{\sqrt{N}} \sum_k S(k) c_{k\uparrow} c_{-k+q\downarrow} \quad (14)$$

we get

$$\langle \{A, A^+\} \rangle \geq \frac{2k_B T}{4 t n} \frac{|\Delta_+ \Omega(q)|^2}{\{(1 - \cos q_x) + (1 - \cos q_y) + \eta(1 - \cos q_z)\}} \quad (15)$$

with $\Omega(q) = \sum_k S(k+q) \langle c_{k\uparrow} c_{-k\downarrow} \rangle / N$. We notice that $\Omega(q) \rightarrow \Delta$ when $q \rightarrow 0$.

In the other hand, we have

$$\langle \{A, A^+\} \rangle = F(q) + R(q)$$

with

$$F(q) = \frac{2}{N} \sum_{k,k'} S(k) S^*(k') \langle c_{k\uparrow} c_{-k+q\downarrow} c_{-k'+q\downarrow}^+ c_{k'\uparrow}^+ \rangle \quad (16)$$

and

$$R(q) = -\frac{1}{N} \sum_k |S(k+q)|^2 \{\langle n_{k\uparrow} \rangle + \langle n_{-k\downarrow} \rangle\} + \frac{1}{N} \sum_k |S(k)|^2 \quad (17)$$

$R(q)$ is a regular function of q .

It is easy to see that

$$\langle c_{k\uparrow}^+ c_{-k+q\downarrow}^+ c_{-k'+q\downarrow} c_{k'\uparrow} \rangle \leq 1 \quad (18)$$

and we deduce that

$$\frac{1}{N} \sum_{q \neq 0} F(q) < \infty \quad (19)$$

where we have use Eq. (7). The inequality (15) becomes

$$\frac{1}{N} \sum_{q \neq 0} F(q) \geq \frac{k_B T}{2nt} \frac{1}{N} \sum_{q \neq 0} \frac{|\Delta_+ \Omega(q)|^2}{\{(1 - \cos q_x) + (1 - \cos q_y) + \eta(1 - \cos q_z)\}} - \frac{1}{N} \sum_{q \neq 0} R(q) \quad (20)$$

Since $\Omega(q)$ and $R(q)$ are regular in q , at small q Eq. (20) becomes

$$\frac{1}{N} \sum_{q \neq 0} F(q) \geq \frac{4k_B T \Delta^2}{nt} \frac{1}{N} \sum_{q \neq 0} \frac{1}{q_x^2 + q_y^2 + \eta q_z^2} - \sum_{q \neq 0} R(q) \quad (21)$$

If now, we take the infinite volume limit for a layered system (with $2L + 1$ planes and with periodic boundary conditions), we will have

$$\frac{1}{2L+1} \sum_{\substack{n_z=-L \\ n_z \neq 0}}^L \frac{1}{(2\pi)^2} \int dq_x dq_y F(q) \geq \frac{4k_B T \Delta^2}{nt} \times$$

$$\frac{1}{2L+1} \sum_{\substack{n_z=-L \\ n_z \neq 0}}^L \frac{1}{(2\pi)^2} \int dq_x dq_y \frac{1}{q_x^2 + q_y^2 + \eta \left(\frac{2\pi n_z}{2L+1} \right)^2} - \quad (22)$$

$$\frac{1}{2L+1} \sum_{\substack{n_z=-L \\ n_z \neq 0}}^L \frac{1}{(2\pi)^2} \int dq_x dq_y R(q) \quad (23)$$

It is clear that in strictly two dimensional Hubbard model ($L = 0$ and $\eta = 0$) the righthandside of Eq. (23) diverges and this in contradiction with Eq. (19) unless $\Delta = 0$ (of course for the case $T \neq 0$). Since $S(k)$ is arbitrary this means that we cannot have $\langle c_{k\uparrow} c_{-k\downarrow} \rangle \neq 0$ for any k .

In the other hand, for quasi-two dimensional case (even for 3 layers : $L = 1$), the Hohenberg proof (Eq. (23)) cannot exclude to have $\langle c_{k\uparrow} c_{-k\downarrow} \rangle \neq 0$.

Then if someone believes that high T_c superconductivity is related to average like $\langle c_{k\uparrow} c_{-k\downarrow} \rangle$, he should keep in mind that he is dealing with a quasi-two-dimensional system and not a strictly two-dimensional one.

In the case of the reduced BCS Hamiltonian quasi-averages $\langle c_{k\uparrow} c_{-k\downarrow} \rangle$ exist even in 1 and 2 dimensions, because the interaction in this Hamiltonian is non local (it depends on quasi-particles velocity) and then violates the f -sum rule (which expresses the particle number conservation) [5, 6]. In practice an other term coming from the interaction is added to the Eq. (12).

1.2 Hubbard model of bosons

The analogous situation holds also for the case of bosonic systems. The extention of our investigation of LRO to Hubbard model for bosons is also motivated by HT_c superconductivity. Especially the question of the existence or not of a bose condensation for HT_c oxides [7].

Let us consider the following Hamiltonian

$$H = \sum_{ij} t_{ij} b_i^\dagger b_j + \sum_{ij} U_{ij} N_i N_j \quad (24)$$

where $(b_i, b_i^\dagger)$ is a boson field, $N_i = b_i^\dagger b_i$ and t_{ij} is given by Eq. (3).

We define the charge density as

$$\rho_i = b_i^+ b_i \quad (25)$$

calculation leads to

$$\begin{aligned} [\rho_q, H], \rho_q^+ &= 4t \sum_k \{ (1 - \cos q_x) \cos k_x + \\ & (1 - \cos q_y) \cos k_y + \eta(1 - \cos q_z) \cos k_z \} N_k \end{aligned} \quad (26)$$

and the following inequality holds

$$0 \langle [\rho_q, H], \rho_q^+ \rangle \leq 4tN \{ (1 - \cos q_x) + (1 - \cos q_z) + \eta(1 - \cos q_z) \} \quad (27)$$

If one uses the Bogoliubov inequality Eq. (8) for

$$C = \rho_q \quad \text{and} \quad A = b_q^+ \quad (28)$$

he gets the following result

$$\frac{1}{N} \sum_{q \neq 0} \langle N_q \rangle \geq -\frac{1}{2} + \frac{k_B T | \langle b_i^+ \rangle |^2}{4t} \frac{1}{N} \sum_{q \neq 0} \frac{1}{(1 - \cos q_x) + (1 - \cos q_y) + \eta(1 - \cos q_z)} \quad (29)$$

Clearly we obtain the same result as for the Fermi systems. In strictly two-dimensional case $\sum_{q \neq 0} \langle N_q \rangle / N$ will be not bounded unless $\langle b_i^+ \rangle = 0$, which implies that there is no LRO at $T \neq 0$. In quasi-two-dimensional case the inequality (29) is not in contradiction with having $\langle b_i^+ \rangle \neq 0$ even at $T \neq 0$.

Finally we have proved that for the Hubbard model (of fermions or bosons), the inequality of Bogoliubov which excludes to have LRO in the two-dimensional case at $T \neq 0$, does not exclude to have LRO for quasi-two-dimensional case even for a finite number of layers. We believe that writing Eliashberg equations for models devoted to describe HT_c superconductivity will be meaningful only if we consider quasi-two-dimensional systems.

References

- [1] P. Monthoux and D. Pines, Phys. Rev. B **47**, 6069 (1993).
- [2] Rice *et al.* preprint, (1994).
- [3] P.C. Hohenberg, Phys. Rev. **158**, 383 (1967).
- [4] T. Hotta, J. Phys. Soc. Jpn. **62**, 4414 (1993).

- [5] P.W. Anderson, Phys. Rev. **110**, 827 (1958).
- [6] N.N. Bogoliubov, D.N. Zubarev and Yu.A. Tserkovnikov, Soviet Phys. - JETP **12**, 88 (1961).
- [7] R. Micnas, J. Ranninger and S. Robaszkiewicz, Rev. Mod. Phys. **62**, N°1, 113 (1990).